

Experimental research basins in Italy: an inventory

Luca BROCCA

Research Institute for Geo-Hydrological Protection, National Research Council, Italy, Perugia

luca.brocca@irpi.cnr.it

The research activity on small experimental basins in Italy started in '70s mainly focusing on runoff generation processes and the hydrologic response time of the catchments. Since then, many isolated initiatives have been carried out expanding significantly research objectives activities.

In this first inventory of the Italian experimental research basins, the information for nine catchments, for which the experimental activity is still ongoing, has been collected and synthesized. Figure 1 shows the geographical location of the basins while Table 1 summarizes their main characteristics. As it can be seen, basins are mainly located in North (3) and South (5) Italy, elevations range between 60 and 3152 m a.s.l. and drainage areas between 1.1 and 72 km². The monitoring of rainfall and runoff is available for all the catchments highlighting that the knowledge of the main components of the water balance is fundamental for more detailed analyses. Additional measurements of several meteorological quantities (temperature, wind, air humidity, solar radiation) along with soil moisture, groundwater level and isotopes are used for improving the understanding on the runoff generation processes. Other basins are equipped for the monitoring of sediment yield and concentrations, bed load transport, nutrients and water quality parameters. For most basins, the monitoring started after 2000 even though a data set of more than 20 years is available for the Gallina and Margiogrande basin.





Figure 1 - Location of small experimental catchments in Italy.

For three catchments (Fiumarella, Vallaccia e Vauz), the research activities are addressed to study: (i) the runoff generation mechanisms for improving flood predictions (Brocca *et al.*, 2009; 2011; Onorati *et al.*, 2009; Penna *et al.*, 2011), (ii) the spatial and temporal variability of soil moisture (Penna *et al.*, 2009; 2012; Brocca *et al.*, 2010b), and (iii) the use of remote sensing for soil moisture retrieval (Brocca *et al.*, 2010a; Manfreda *et al.*, 2011). These activities are carried out by using different rainfall-runoff models characterized by different complexity and spatial discretization. For two basins, several models are applied to reproduce not only the rainfall-runoff process but also water quality parameters and nutrient concentrations (Celone, De Girolamo *et al.*, 2011) and soil erosion (Cannata, Licciardello *et al.*, 2007; 2011). Sediment and bed load transport is investigated at the Gallina, Margiogrande and Rio Cordon (Lenzi *et al.*, 2004) basins. Specifically, long-term evaluation is

ERB Newsletter 2013 - 1



carried out at Margiogrando (20 years) and Gallina (30 years) basins also to understand the possible impact of climate changing on this process (Anselmo *et al.*, 2011). The estimation of river velocity and flow resistance for stepped channels is also analyzed at Rio Cordon (Comiti *et al.*, 2007). Finally, the impact of check-dams on fluvial landforms and riparian vegetation is the main research activity at Sant'Agata basin (Bombino *et al.*, 2006; 2009).

As it can be inferred from the brief summary reported above, investigation in small experimental basins in Italy have provided very significant data sets that allowed a better understanding of several scientific issues. This also resulted in the publication of several papers on major international scientific journals. Moreover, to our knowledge, several other small research basins are present in Italy that have not yet been included in the ERB network. It is our goal to stimulate also their participation. Future steps of this inventory will be the formation of a strong network for the harmonization of the research activities in the different experimental basins also by sharing the collected data sets. This will permit researchers to analyze a more extended database (both in time and in space) thus obtaining more robust results. Therefore, a significant advance in our understanding of the hydrological cycle is expected.


Table 1 - Main characteristics of the Italian experimental catchments (R: rainfall, T: temperature, H: air humidity, W: wind velocity/direction, Rad: solar radiation, Q: discharge, SM: soil moisture, G: groundwater level, SED: sediments concentration/yield, BL: bed load, S: snow, Nut: nutrient concentration, WQ: water quality).

Name	Area (km ²)	Lat/Lon (°)	Elevation range (m a.s.l.)	Mean annual R (mm) / T (°C)	Collected data	Start year	Main contact
Cannata	1.3	37°53'N/ 14°46'E	903-1270	690 / NA	R, Q, SED	1996	flicciar@uniict.it
Celone	72.0	41°23'N/ 15°20'E	60-1150	730 / 13	R, T, Q, Nut	2009	degirolamo@ba.irsa.cnr.it
Fiumarella	32.5	40°24'N/ 16°03'E	630-1380	600 / 10	R, T, H, W, S, Rad, Q, SM	2002	salvatore.manfreda@unibas.it
Gallina	1.1	45°38'N/ 8°18'E	330-522	1280 / 11	R, T, Q, SED	1982	franca.maraga@gmail.com
Margiogrando	4.5	37°48'N/ 14°52'E	580-1030	600 / NA	R, Q, GE	1993	dzema@unirc.it
Rio Cordon	5.0	46°27'N/ 12°06'E	1763-2748	1100 / 2	R, T, H, Rad, S, Q, SED, BL, WQ	1985	marioaristidelenzi@unipd.it
Sant'Agata	61.0	38°08'N/ 15°53'E	893-1610	1350 / 11	R, Q	2003	giuseppe.bombino@unirc.it
Vallaccia	57.6	43°16'N/ 12°12'E	288-818	930 / 13	R, T, H, W, Q, SM	1988	luca.brocca@irpi.cnr.it
Vauz	1.9	46°29'N/ 11°50'E	1847-3152	1220 / 3	R, T, Q, SM, G, I	2005	daniele.penna@unipd.it

References

- Anselmo, V., Caroni, E., Maraga, F., Pelissero, C. (2011). Tendencies in the hydrological budget of a pre-alpine small basin. *Journal for Land Management, Food and Environment*, 62, 9-15.
- Bombino, G., Tamburino, V., Zimbone, S.M. (2006). Assessment of the effects of check-dams on riparian vegetation in the mediterranean environment: a methodological approach and applications. *Ecological Engineering*, 27, 134-144.
- Bombino, G., Gurnell, A.M., Tamburino, V., Zema, D.A., Zimbone, S.M. (2009). Adjustments in channel form, sediment calibre and vegetation around check dams in the headwater reaches of mountain torrents, Calabria, Italy. *Earth Surface Processes and Landforms*, 34, 1011-1021.
- Brocca, L., Melone, F., Moramarco, T., Singh, V.P. (2009). Assimilation of observed soil moisture data in storm rainfall-runoff modelling. *Journal of Hydrologic Engineering*, 14 (2), 153-165.
- Brocca, L., Melone, F., Moramarco, T., Wagner, W., Hasenauer, S. (2010a). ASCAT Soil Wetness Index validation through in-situ and modeled soil moisture data in central Italy. *Remote Sensing of Environment*, 114 (11), 2745-2755.
- Brocca, L., Melone, F., Moramarco, T., Morbidelli, R. (2010b). Spatial-temporal variability of soil moisture and its estimation across scales. *Water Resources Research*, 46, W02516.

ERB Newsletter 2013 - 1



- Brocca, L., Melone, F., Moramarco, T., Penna, D., Borga, M., Matgen, P., Heitz, S. (2011). Investigation of the hydrologic response of three experimental basins across Europe. *Journal for Land Management, Food and Environment*, 62, 31-37.
- Comiti F., Mao L., Wilcox A., Wohl E., Lenzi M.A. (2007). Field-derived relationships for flow velocity and resistance in high-gradient streams. *J Hydrol.*, 340, 48-62.
- De Girolamo, A.M., Calabrese, A., Lo Porto, A., Oueslati, O., Pappagallo, G., Santese, G. (2011). Hydrologic regime characterization for a semi-arid watershed. *Journal for Land Management, Food and Environment*, 62, 39-45.
- Lenzi, M.A., Mao, L., and Comiti, F. (2004). Magnitude-frequency analysis of bed load data in an Alpine boulder bed stream. *Water Resources Research*, 40, W07201.
- Licciardello, F., Zema, D.A., Zimbone, S.M., Bingner, R.L. (2007). Runoff and soil erosion evaluation by the AnnAGNPS model in a small Mediterranean watershed. *Transactions of the ASABE*, 50(5), 1585-1593.
- Licciardello, F., Rossi, C.G., Srinivasan, R., Zimbone, S.M., Barbagallo, S. (2011). Hydrologic evaluation of a Mediterranean watershed using the SWAT model with multiple pet estimation methods. *Transactions of the ASABE*, 54(5), 1615-1625.
- Manfreda, S., Lacava, T., Onorati, B., Pergola, N., Di Leo, M., Margiotta, M.R., Tramutoli, V. (2011). On the use of AMSU-based products for the description of soil water content at basin scale. *Hydrol. Earth Syst. Sciences*, 15, 2839-2852.
- Onorati, B., Margiotta, M.R., Carriero, D., Manfreda, S., Fiorentino, M. (2009). Experimental Evidence on Runoff Generation Mechanisms, Variability in space and time of extreme rainfalls, floods and droughts. *Proceedings of the AMHY-FRIEND International Workshop on Hydrological Extremes*, Cosenza (Italy), June 6-8.
- Penna, D., Borga, M., Norbiato, D., Dalla Fontana, G. (2009). Hillslope scale soil moisture variability in a steep alpine terrain. *J Hydrol.*, 364, 311-327.
- Penna, D., Tromp-Van Meerveld, H. J., Gobbi, A., Borga, M., Dalla Fontana, G. (2011). The influence of soil moisture on threshold runoff generation processes in an alpine headwater catchment. *Hydrol. Earth Syst. Sciences*, 15, 689-702.
- Penna, D., Brocca, L., Borga, M., Dalla Fontana, G. (2012). Soil moisture temporal stability on two alpine hillslopes during wet and dry periods. *J Hydrol.*, in press.